

Model based approach for household clustering with mixed scale variables

CHRISTIAN CARMONA¹, LUIS NIETO-BARAJAS² & ANTONIO CANALE³

¹ *Department of Statistics, University of Oxford*

² *Department of Statistics, ITAM*

³ *Department of Economics and Statistics, University of Turin*

carmona@stats.ox.ac.uk, lnieto@itam.mx and antonio.canale@unito.it

Abstract

The Ministry of Social Development in Mexico is in charge of creating and assigning social programmes targeting specific needs in the population for the improvement of quality of life. To better target the social programmes, the Ministry is aimed to find clusters of households with the same needs based on demographic characteristics as well as poverty conditions of the household. Available data consists of continuous, ordinal, and nominal variables and the observations are not iid but come from a survey sample based on a complex design. We propose a Bayesian nonparametric mixture model that jointly models this mixed scale data and accommodates for the different sampling probabilities. The performance of the model is assessed via simulated data. A full analysis of socio-economic conditions in households in the State of Mexico is presented.

Keywords: Bayes nonparametrics, complex design, latent variables, multivariate normal, Poisson-Dirichlet process.

1 Introduction

The Ministry of Social Development (SEDESOL, according to its initials in Spanish) in Mexico is one of the many government dependencies. The aim of SEDESOL is to help and improve the social backwardness that prevails in a high percentage of the households in

the country. To fulfil this objective, SEDESOL creates social programmes to target specific needs in the population.

Currently, each existing social programme has its own rules of operation and its own means of selecting the potential households to be benefited by the programme, but in general they all use the income as the main selection criterion. In order to simplify the selection of potential candidates and to better target the programmes to the correct population, SEDESOL wants to create a clustering of households based on needs, socio-economic and demographic characteristics as well as poverty conditions.

In 2009, the Council for National Evaluation (Coneval, 2009) proposed a methodology for measuring the poverty conditions in households in terms of multiple indicators. These include the income dimension, social deprivations and social cohesion. As a result of this new methodology a multi-dimensional measurement was created based on seven indicators: income, and six deprivation indicators such as education, access to health services, access to social security, housing quality, access to basic public services and access to feeding. These will be the core set of variables used in the clustering later on.

In January of 2004 a new general law for social development was promulgated in Mexico. It establishes that poverty measurements must be done every two years at a state level and every five years at a municipality level. For these purposes, the National Institute for Official Statistics (INEGI) implements a survey based on a complex design of households. This national survey of income and expenses in households (ENIGH) through a module of socio-economic conditions (MCS) collects the necessary information to produce the multi-dimensional poverty indicators at a household level and in some cases at an individual level. These household poverty indicators are then expanded with the corresponding sampling design weights to produce poverty indicators at a state level.

The aim of this work is to produce clusters of households based primarily on the multidimensional poverty indicators considering that the available data come from a complex

sampling design where each household has a different sampling probability. As mentioned, the data is a result of mixed scale variables that include continuous, discrete, ordinal and nominal variables. We introduce a convenient set of latent variables associated to each of the observed ones and use a Bayesian nonparametric mixture of multivariate normals to model their latent multivariate density. The model also accounts for the different sampling probabilities of the selected households.

Model based clustering approaches (Banfield and Raftery, 1993) rely on a probability model based on a finite or infinite mixture of sub-populations. Typically the sub-populations are assumed to be normal but other alternatives, to deal for example with possible asymmetric structures of the subpopulation, have also been considered (Rodríguez and Walker, 2014; Canale and Scarpa, 2016). In the case of mixed-scale data, Everitt (1988) uses latent variables and thresholding to deal with binary and ordinal variables. Although he did not performed an actual clustering, the author mentions that his maximum likelihood estimation procedure can be potentially used for clustering. Many authors have considered variations of the use of latent variable and thresholding to induce a multivariate mixed-scale density such as Norets and Pelenis (2012), McParland et al. (2014) or Canale and Dunson (2015). Specifically, McParland et al. (2014) propose a finite mixture model to accomplish a similar task as the one considered here. As in our motivating application, their setting consists of mixed-scale data, and specifically of binary, ordinal and nominal variables. To accommodate the nominal variables, the authors propose an extended version of the latent variables considered in Everitt (1988). They further parametrized the mean of the latents via a factor model, reducing so the effective number of parameters to estimate but losing some information. Our approach is similar to McParland et al. (2014) but differs in several ways: we include continuous variables to the mixed-mode set of variables, we propose an infinite mixture, we do not reduce dimensionality in the parametrization of the mean of the latent variables. In addition we account for the complex design setting of our data while the

paper by McParland et al. (2014) focuses only on iid data.

The layout of the remaining of the paper is as follows: in Section 2 we define the model and the prior used to induce the clustering. Section 3 contains posterior characterisation of the model parameters, with some implementation details included in Section 4. A simulation study and the analysis of the motivating data set are presented in Section 5. We end with some concluding remarks in Section 6.

Before we proceed we introduce notation: $N_q(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ denotes a q -variate normal distribution with mean vector $\boldsymbol{\mu}$ and variance-covariance matrix $\boldsymbol{\Sigma}$; $\text{Be}(a, b)$ denotes a beta distribution with mean $a/(a + b)$; $\text{Ga}(a, b)$ denotes a gamma distribution with mean a/b ; $\text{IGa}(a, b)$ denotes an inverse gamma distribution with mean $b/(a - 1)$.

2 Model

2.1 Observed and latent variables

Consider an individual i that is characterized by a multivariate response of dimension p , i.e., $\mathbf{y}_i = \{y_{ij}, j = 1 \dots, p\}$ and $i = 1, \dots, n$. Let us assume that the total number of variables p is divided into c continuous variables, o ordinal variables, and m nominal variables such that $p = c + o + m$. In summary, $\mathbf{y}'_i = (y_{i,1}, \dots, y_{i,c}, y_{i,c+1}, \dots, y_{i,c+o}, y_{i,c+o+1}, \dots, y_{i,c+o+m})$ for $i = 1 \dots, n$.

We associate to each response $\mathbf{y}_i$ of dimension p a latent vector $\mathbf{z}_i$ of dimension q in the following way:

- For each *continuous* variable y_{ij} , $j = 1, \dots, c$, we define a latent $z_{ij} = g_j(y_{ij})$, where $g_j(\cdot)$ is a normalising transformation possibly different for each j .
- For each *ordinal* variable y_{ij} , $j = c + 1, \dots, c + o$, that takes values in an ordered set $\{\vartheta_k\}$ with K_j different values, we need one latent z_{ij} that satisfies $y_{ij} = \vartheta_k$ iff $\gamma_{j,k-1} < z_{ij} \leq \gamma_{j,k}$, for $k = 1, \dots, K_j$ and $\{\gamma_{j,0}, \dots, \gamma_{j,K_j}\}$ are fixed thresholds or cut-off

points with $\gamma_{j,0} = -\infty$ and $\gamma_{j,K_j} = \infty$. More details on the choice of the cut-offs are discussed later. Note that a binary variable is a special case of an ordinal variable with $K_j = 2$.

- For each *nominal* variable y_{ij} , $j = c + o + 1, \dots, c + o + m$, with L_j categories we need $L_j - 1$ latent variables that can be placed in the vector $\mathbf{z}_{ij} = \{z_{il}, l = c + o + \sum_{h=c+o+1}^{j-1} (L_h - 1) + 1, \dots, c + o + \sum_{h=c+o+1}^j (L_h - 1)\}$ such that

$$y_{ij} = \begin{cases} L_j, & \text{if } \max_l(z_{il}) < 0 \\ k, & \text{if } z_{i,s} = \max_l(z_{il}) \text{ \& } z_{i,s} > 0 \\ & \text{with } s = c + o + \sum_{h=c+o+1}^{j-1} (L_h - 1) + k, \text{ and } k = 1, \dots, L_j - 1 \end{cases}$$

where $q = c + o + \sum_{j=c+o+1}^{c+o+m} (L_j - 1)$. In summary $\mathbf{z}'_i = (z_{i,1}, \dots, z_{i,c}, z_{i,c+1}, \dots, z_{i,c+o}, \mathbf{z}_{i,c+o+1}, \dots, \mathbf{z}_{i,c+o+m})$ which for $j = c + o + 1, \dots, c + o + m$ each $\mathbf{z}_{i,j}$ is of dimension $L_j - 1$.

Available data come from a complex survey sample where each individual $\mathbf{y}_i$ has known sampling probability π_i , $i = 1, \dots, n$. The reciprocal of these sampling probabilities, $w_i = 1/\pi_i$, are called sampling design weights or expansion factors.

There are different strategies to include the sample probabilities into an inferential procedure. In linear regression, Lumley (2010) proposes a weighted least squares with weights defined by w_i , that is $\min \sum_{i=1}^n \frac{1}{\pi_i} (y_i - \alpha - \beta x_i)^2$, where x_i is an explanatory variable. Chambers and Skinner (2003), on the other hand, propose to re-weight the likelihood contribution with an exponent given by the sampling weight w_i , i.e., $\prod_{i=1}^n f(y_i | \theta)^{1/\pi_i}$. Considering Lumley's approach, the weighted least squares estimates are equivalent to the maximum likelihood estimators under a normal model with variances scaled by a factor π_i . Moreover, under a normal model Chambers and Skinner's approach is also equivalent to scale the variance with a factor π_i . Therefore we propose the following model:

$$\mathbf{z}_i | \boldsymbol{\mu}_i, \boldsymbol{\Sigma} \sim N_q(\boldsymbol{\mu}_i, \kappa \pi_i \boldsymbol{\Sigma}), \quad (1)$$

where $\boldsymbol{\mu}_i$ is a mean vector of dimension $q \times 1$, $\boldsymbol{\Sigma}$ is a variance-covariance matrix of dimension $q \times q$ and κ is a scaling factor that controls the impact of the sampling probabilities in the

variance. This latter parameter will play a central role in the posterior clustering structures as discussed later in Section 5.

Since there is a deterministic relationship between $\mathbf{y}_i$ and $\mathbf{z}_i$, and some of the variables in $\mathbf{y}_i$ are categorical (ordinal or nominal), some estimability constraints have to be imposed in the matrix $\mathbf{\Sigma}$. If we denote by σ_j^2 the variance of the j th variable and by $\sigma_{j,k}$ the covariance between variables j and k for $j, k = 1, \dots, n$, then:

- (i) For $j = 1, \dots, c : \sigma_j^2 > 0$,
- (ii) For $j = c + 1, \dots, c + o : \sigma_j^2 = 1$
- (iii) For $j = c + o + 1, \dots, \sum_{h=c+o+1}^{c+o+m} (L_h - 1) : \sigma_j^2 = 1$.

This last specification, along with the mixture model carefully described in the next section, is crucial as it implies that the internal cut-offs for the ordinal variables can be fixed without loss of generality.

2.2 Prior distributions

The clustering model will be based in an appropriate choice of the prior distribution on the $\boldsymbol{\mu}_i$'s. A clustering of the $\boldsymbol{\mu}_i$'s will induce a clustering of the $\mathbf{y}_i$'s. For that we want the marginal distribution for each $\boldsymbol{\mu}_i$ to have a continuous support on the real line and the joint distribution to satisfy that $\Pr(\boldsymbol{\mu}_i = \boldsymbol{\mu}_{i'}) > 0$ for all $i \neq i' \in \{1, \dots, n\}$. There are several nonparametric priors that satisfy these conditions. A sufficiently rich class is the (two-parameters) Poisson-Dirichlet process (Pitman and Yor, 1997) that includes the Dirichlet (Ferguson, 1973) and the normalized stable (Kingman, 1975) processes as particular cases.

A Poisson-Dirichlet process G is almost surely discrete (see e.g. Ishwaran and James, 2001) and is defined as follows:

$$G(\cdot) = \sum_{k=1}^{\infty} w_k \delta_{\xi_k}(\cdot),$$

where δ_ξ denotes a point mass at ξ . Here, both the weights w_k and the locations ξ_k are random variables such that $\xi_k \stackrel{\text{iid}}{\sim} G_0$ for all $k = 1, 2, \dots$, with G_0 a probability distribution. The heights $\{w_k\}$ are defined as $w_1 = v_1$ and $w_k = v_k \prod_{l < k} (1 - v_l)$, with $v_k \stackrel{\text{ind}}{\sim} \text{Be}(1 - a, b + ka)$. This specific choice for the distribution of the stick-breaks v_k characterizes the Poisson-Dirichlet process. The parameter G_0 is known as centering measure since $E(G) = G_0$. In notation we say $G \sim \mathcal{PD}(a, b, G_0)$ with parameters $a \in [0, 1)$, $b > -a$ and centring measure G_0 . The Dirichlet and the normalized stable processes arise when $a = 0$ and when $b = 0$, respectively.

Therefore our prior on the $\boldsymbol{\mu}_i$'s will be

$$\boldsymbol{\mu}_i | G \stackrel{\text{iid}}{\sim} G, \text{ for } i = 1, \dots, n \text{ with } G \sim \mathcal{PD}(a, b, G_0), \quad (2)$$

and $G_0(\boldsymbol{\mu}) = N(\mathbf{0}, \boldsymbol{\Sigma}_\mu)$, where $\boldsymbol{\Sigma}_\mu = \text{diag}(\sigma_{\mu 1}^2, \dots, \sigma_{\mu q}^2)$. In consequence, this choice of prior implies that the $\boldsymbol{\mu}_i$'s are exchangeable with marginal distribution $\boldsymbol{\mu}_i \sim G_0$ for all $i = 1, \dots, n$. Pitman (1995) showed that if we integrate out the nonparametric measure G , the joint distribution of the $\boldsymbol{\mu}_i$'s is characterized by a generalized Polya urn mechanism with conditional distribution that depends on the density g_0 associated to G_0 and given by

$$f(\boldsymbol{\mu}_i | \boldsymbol{\mu}_{-i}) = \frac{b + a r_i}{b + n - 1} g_0(\boldsymbol{\mu}_i) + \sum_{j=1}^{r_i} \frac{n_{j,i}^* - a}{b + n - 1} \delta_{\boldsymbol{\mu}_{j,i}^*}(\boldsymbol{\mu}_i), \quad (3)$$

for $i = 1, \dots, n$, where $\boldsymbol{\mu}_{-i} = (\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_{i-1}, \boldsymbol{\mu}_{i+1}, \dots, \boldsymbol{\mu}_n)$ denotes the set of all $\boldsymbol{\mu}_j$'s excluding the i^{th} element, and $(\boldsymbol{\mu}_{1,i}^*, \dots, \boldsymbol{\mu}_{r_i,i}^*)$ denote the unique values in $\boldsymbol{\mu}_{-i}$, each occurring with frequency $n_{j,i}^*$, $j = 1, \dots, r_i$, which satisfy the condition $n_{1,i}^* + \dots + n_{r_i,i}^* = n - 1$. Therefore $\Pr(\boldsymbol{\mu}_i = \boldsymbol{\mu}_{i'}) > 0$ as desired. The number of unique values r in $\boldsymbol{\mu} = (\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_n)$ determines the number of clusters. This value is controlled by the parameters (a, b) . Larger values of either a or b , within the valid ranges, produce a larger r (e.g. Navarrete et al., 2008).

The mixture specification allows the threshold for the ordinal variables to be fixed without loss of generality as discussed in Kottas et al. (2005), Leon-Novelo et al. (2010), and

Bandyopadhyay and Canale (2016). Indeed, without the mixture specification and assuming a single multivariate normal density for the latent variables, the model can be thought as a multivariate ordered probit models. In this models, for the sake of identifiability, each first marginal cut-off is fixed, and the conditional means are free to vary. However, the nonparametric mixture model provides enough flexibility to fix the cut-offs while placing different kernel densities within each interval so to have varying probability mass assigned to each ordinal category. Specifically, given the fixed variance specification (ii) and (iii), the internal thresholds can be fixed at $\gamma_{j,k} \in \{0, \pm 4, \pm 8, \dots\}$ as discussed in Leon-Novelo et al. (2010).

Considering the variance-covariance matrix Σ we recall that it has several constraints on the diagonal given by conditions (i)–(iii). To include them and to ensure positive definiteness, we follow a separation strategy as in Barnard et al. (2000) such that $\Sigma = \Lambda \Omega \Lambda$ where $\Lambda = \text{diag}(\sigma_1, \dots, \sigma_q)$ is a diagonal matrix of standard deviations and Ω is a correlation matrix. We thus assign independent priors on the squared elements of Λ and on Ω .

For those $j = 1, \dots, q$ such that $\sigma_j^2 \neq 1$ we take

$$\sigma_j^2 \stackrel{\text{iid}}{\sim} \text{IGa}(d_0^z, d_1^z), \quad (4)$$

and for the correlation matrix we take

$$f(\Omega) \propto |\Omega|^{q(q-1)/2-1} \left(\prod_j |\Omega_{jj}| \right)^{-(q+1)/2}, \quad (5)$$

where Ω_{jj} is the i th principal sub-matrix of Ω . This prior implies that individual correlations are marginally uniform on the interval $[-1, 1]$ (Barnard et al., 2000).

We finally take hyper-priors for the parameters a , b and $\sigma_{\mu j}$ as

$$\begin{aligned} f(a) &= \alpha \delta_0 + (1 - \alpha) \text{Be}(a \mid d_0^a, d_1^a), \quad f(b \mid a) = \text{Ga}(b + a \mid d_0^b, d_1^b), \\ \text{and } \sigma_{\mu j}^2 &\stackrel{\text{iid}}{\sim} \text{IGa}(d_0^\mu, d_1^\mu) \quad j = 1, \dots, q. \end{aligned} \quad (6)$$

Note that the hyper-prior on a is a mixture of a point mass at zero and a continuous beta distribution. This is to consider the option that $a = 0$ with positive probability. The hyper

prior on b is given conditionally on a and includes the constraint that $b > -a$ by shifting the support of the gamma density to the interval $(-a, \infty)$. The values $d_0^z, d_1^z, d_0^a, d_1^a, d_0^b, d_1^b, d_0^\mu, d_1^\mu$ are all known constants.

3 Posterior characterisation

Let $(\boldsymbol{\mu}, \boldsymbol{\Sigma}, a, b, \boldsymbol{\Sigma}_\mu)$ the set of parameters and hyper-parameters of the model, where $\boldsymbol{\mu}' = (\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_n)$. The posterior distribution of these parameters is characterised in terms of its full conditional distributions, given below:

- (a) The conditional posterior distribution of $\boldsymbol{\mu}_i$, $i = 1, \dots, n$, is

$$f(\boldsymbol{\mu}_i \mid \mathbf{z}, \boldsymbol{\mu}_{-i}) = p_0 N_q(\boldsymbol{\mu}_i \mid \boldsymbol{\nu}_i, \mathbf{V}_i) + \sum_{j=1}^{r_i} p_j \delta_{\boldsymbol{\mu}_{j,i}^*}(\boldsymbol{\mu}_i),$$

where $\boldsymbol{\nu}_i = \mathbf{V}_i(\pi_i \kappa \boldsymbol{\Sigma})^{-1} \mathbf{z}_i$, $\mathbf{V}_i = ((\pi_i \kappa \boldsymbol{\Sigma})^{-1} + \boldsymbol{\Sigma}_\mu^{-1})^{-1}$, $p_j = D_j / (\sum_{l=0}^{r_i} D_l)$ for $j = 0, \dots, r_i$ with $D_0 = (b + ar_i) N_q(\mathbf{z}_i \mid \mathbf{0}, \pi_i \kappa \boldsymbol{\Sigma} + \boldsymbol{\Sigma}_\mu)$ and, for $j > 0$, $D_j = (n_{ji}^* - a) N_q(\mathbf{z}_i \mid \boldsymbol{\mu}_{ji}^*, \pi_i \kappa \boldsymbol{\Sigma})$.

Conditional distribution (a) allows us to identify which μ_i 's are equal to each other or different. We further need to re-sample the unique values $(\boldsymbol{\mu}_i^*, \dots, \boldsymbol{\mu}_r^*)$ in $(\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_n)$.

- (b) Conditional on the membership allocations $I_j = \{i : \boldsymbol{\mu}_i = \boldsymbol{\mu}_j^*\}$, $j = 1, \dots, r$

$$f(\boldsymbol{\mu}_j^* \mid \mathbf{z}, I_j, \text{rest}) = N(\boldsymbol{\mu}_j^* \mid \boldsymbol{\nu}_j^*, \mathbf{V}_j^*),$$

where $\boldsymbol{\nu}_j^* = \frac{1}{\kappa} \mathbf{V}_j^* \boldsymbol{\Sigma}^{-1} \left(\sum_{i \in I_j} (1/\pi_i) \mathbf{z}_i \right)$ and $\mathbf{V}_j^* = \left(\left(\frac{1}{\kappa} \sum_{i \in I_j} (1/\pi_i) \right) \boldsymbol{\Sigma}^{-1} + \boldsymbol{\Sigma}_\mu^{-1} \right)^{-1}$.

- (c) The conditional posterior distribution of the diagonal elements $\sigma_{\mu l}^2$, $l = 1, \dots, q$, of matrix $\boldsymbol{\Sigma}_\mu$ is

$$f(\sigma_{\mu l}^2 \mid \text{rest}) = \text{IGa} \left(\sigma_{\mu l}^2 \mid d_0^\mu + \frac{r}{2}, d_1^\mu + \frac{1}{2} \sum_{j=1}^r (\mu_{jl}^*)^2 \right),$$

where μ_{jl}^* is the l -th coordinate of vector $\boldsymbol{\mu}_j^*$.

For the matrix $\mathbf{\Sigma}$, considering the separation strategy, the full conditional distributions of $\mathbf{\Lambda}$ and $\mathbf{\Omega}$ are

- (d) The conditional posterior distribution of the squared diagonal elements of $\mathbf{\Lambda}$, σ_j^2 , for those j such that $\sigma_j^2 \neq 1$, is given by

$$f(\sigma_j^2 \mid \mathbf{z}, \text{rest}) \propto (\sigma_j^2)^{-(d_0^z + n/2 + 1)} e^{-d_1^z / \sigma_j^2} \exp \left\{ -\frac{1}{2} \text{tr}(\mathbf{\Sigma}^{-1} \mathbf{S}) \right\},$$

where $\mathbf{S} = \sum_{i=1}^n \frac{1}{\kappa \pi_i} (\mathbf{z}_i - \boldsymbol{\mu}_i)(\mathbf{z}_i - \boldsymbol{\mu}_i)'$.

- (e) The conditional posterior distribution of $\mathbf{\Omega}$ is simply

$$f(\mathbf{\Omega} \mid \mathbf{z}, \mathbf{\Lambda}, \text{rest}) \propto \left(\prod_{j=1}^q |\mathbf{\Omega}_{jj}| \right)^{-\frac{1}{2}(q+1)} |\mathbf{\Omega}|^{-\frac{1}{2}\{n+2-q(q-1)\}} \exp \left[-\frac{1}{2} \text{tr} \{ \mathbf{\Omega}^{-1} (\mathbf{\Lambda}^{-1} \mathbf{S} \mathbf{\Lambda}^{-1}) \} \right],$$

where $\mathbf{S}$ is given above.

The likelihood for a and b is given by the exchangeable partition probability function induced by the Poisson-Dirichlet process (Pitman, 1995). Thus,

- (f) The conditional posterior distribution of a is

$$f(a \mid b, \text{rest}) \propto \left\{ \prod_{j=1}^{r-1} (b + ja) \right\} \left\{ \prod_{j=1}^r \frac{\Gamma(n_j^* - a)}{\Gamma(1 - a)} \right\} f(a),$$

where the prior $f(a)$ is given in (6).

- (g) The conditional posterior distribution of b is

$$f(b \mid a, \text{rest}) \propto \frac{\Gamma(b+1)}{\Gamma(b+n)} \left\{ \prod_{j=1}^{r-1} (b + ja) \right\} f(b \mid a),$$

where $f(b \mid a)$ is also given in (6).

Finally, the latent variables z_{ij} have to be re-sampled from their respective predictive distributions following the constraints mentioned at the beginning of Section 2.

(h) The conditional predictive distributions for z_{ij} , $i = 1, \dots, n$ are:

- For $j = 1, \dots, c$, z_{ij} remains unchanged.
- For $j = c + 1, \dots, c + o$, z_{ij} has a truncated normal distribution of the form

$$z_{ij} \mid \mathbf{z}_{-(ij)}, \text{rest} \sim N(\nu_{ij}, V_{ij})I(\gamma_{j,k-1} < z_{ij} \leq \gamma_{j,k}),$$

for $y_{ij} = k$, where $\nu_{ij} = \mu_{ij} + \Sigma_{12}\Sigma_{22}^{-1}(\mathbf{z}_{-(ij)} - \boldsymbol{\mu}_{-(ij)})$, $V_{ij} = \Sigma_{11} - \Sigma_{12}\Sigma_{22}^{-1}\Sigma_{21}$, with $\Sigma_{11} = \text{Var}(z_{ij}) = \sigma_j^2$, $\Sigma_{12} = \text{Cov}(z_{ij}, \mathbf{z}_{-(ij)})$, and $\Sigma_{22} = \text{Var}(\mathbf{z}_{-(ij)})$.

- For $j = c + o + 1, \dots, c + o + \sum_{h=c+o+1}^m (L_h - 1)$, the latent vector $\mathbf{z}_{ij} = \{z_{il}\}$, $l = c + o + \sum_{h=c+o+1}^{j-1} (L_h - 1), \dots, c + o + \sum_{h=c+o+1}^j (L_h - 1)$ has z_{il} element coming from a truncated normal distribution of the form

$$z_{il} \mid \mathbf{z}_{il}, \text{rest} \sim N(\nu_{il}, V_{il})I_{A_{ij}}(z_{il}),$$

where

$$A_{ij} = \begin{cases} (-\infty, 0) & \text{if } y_{ij} = L_j \\ (-\infty, z_{is}) & \text{if } y_{ij} = k < L_j \text{ \& } l \neq s = c + o + \sum_{h=c+o+1}^{j-1} (L_h - 1) + k \\ (\max_s \{z_{is}, 0\}, \infty) & \text{if } y_{ij} = k < L_j \text{ \& } s \neq l = c + o + \sum_{h=c+o+1}^{j-1} (L_h - 1) + k \end{cases}$$

and ν_{il} and V_{il} are as above.

4 Implementation details and clustering selection

Sampling from conditional distributions (a)–(c) is straightforward. To sample from conditional distribution (d) we propose a random walk Metropolis-Hastings (MH) step (Tierney, 1994). At iteration $(r+1)$ we sample from the proposal distribution $\sigma_j^{2*} \sim \text{Ga}(\delta_\sigma, \delta_\sigma / \sigma_j^{2(r)})$. We first check positive definiteness of Σ^* with the new draw and accept its corresponding probability

For conditional distribution (e) we proceed to sample conditionally one at a time the elements ρ_{jk} of $\boldsymbol{\Omega}$. Using Barnard et al. (2000)'s ideas, we obtain the support of ρ_{jk} that keeps

Ω positive defined by computing the roots of the quadratic function $h(\rho) = \theta_1\rho^2 + \theta_2\rho + \theta_3$, where $\theta_1 = \{h(1)+h(-1)-2h(0)\}/2$, $\theta_2 = \{h(1)-h(-1)\}/2$ and $\theta_3 = h(0)$ with $h(\rho) = |\Omega(\rho)|$ and $\Omega(\rho)$ is the correlation matrix Ω evaluated in ρ at the entry j, k . We can then use the griddy Gibbs sampler (Ritter and Tanner, 1992) and evaluate the conditional density top sample from ρ_{jk} . However, from our experience, this procedure highly increases the computational time. Instead we implement a random walk MH step with uniform proposal distribution. If we denote by $[\rho_1, \rho_2]$ the support of ρ_{jk} and by $\ell = \rho_2 - \rho_1$ its length, the proposal is taken as $\rho_{jk}^* \sim \text{Un}\left(\max(\rho_1, \rho_{jk}^{(r)} - \ell/\delta_\rho), \min(\rho_2, \rho_{jk}^{(r)} + \ell/\delta_\rho)\right)$.

To sample from conditional distributions (f) we implement a MH step with independent proposal distribution $p(a) = (1/2)I_0(a) + (1/2)\text{Be}(a \mid 1, 1)$. For (g) we propose a random walk MH step with proposal distribution as the one used for ρ_{jk} , that is, $b^* \mid b^{(r)} \sim \text{Un}(b^{(r)} - \delta_b, b^{(r)} + \delta_b)$, always considering the support $b > -a$.

In all cases δ_σ , δ_ρ and δ_b are tuning parameters that control the acceptance probability of the corresponding MH steps. For the examples considered here we took $\delta_\sigma = 10$, $\delta_\rho = 10$ and $\delta_b = 2$.

Summarising the posterior distribution assigned to all possible partitions of the data is not an easy task. This posterior distribution is characterised through a MCMC sample. At each iteration of the MCMC sampler an $n \times n$ adjacency matrix containing a 1 in position ij if elements i and j share the same value of μ^* is stored. At the end of the Gibbs sampler, a similarity matrix is computed as the Montecarlo average of all the adjacency matrices. This similarity matrix represents the “average clustering”. As posterior summary of the clustering structure, we select the adjacency matrix with minimum squared distance from the averaged similarity matrix. This procedure was originally proposed by Dahl (2006) and has been recently formalised in a decision theory framework by Wade and Ghahramani (2015).

This posterior inference procedure and the clustering selection has been implemented in R (R Core Team, 2016) in the package `BNPMIXclust` that is available from The Comprehensive

R Archive Network (CRAN).

5 Data analysis

5.1 Simulated data

In order to evaluate the clustering performance of our model we carried out a simulation study. We sampled 3-dimensional latent continuous vectors $\mathbf{z}' = (z_1, z_2, z_3)$ from a 3-components mixture of normals with equal mixing probabilities and mixture components with means $\boldsymbol{\mu}'_1 = (2, 2, 5)$, $\boldsymbol{\mu}'_2 = (6, 4, 2)$ and $\boldsymbol{\mu}'_3 = (1, 6, 2)$, and variances $\boldsymbol{\Sigma}_1 = \text{diag}(1, 1, 1)$, $\boldsymbol{\Sigma}_2 = \text{diag}(0.1, 2, 0.1)$ and $\boldsymbol{\Sigma}_3 = \text{diag}(2, 0.1, 0.1)$. A sample from this model is included in Figure 1. The data clearly show the existence of three groups.

We tested the model by considering different scenarios for the observed data that combine the use continuous and discretised versions of the simulated data. These scenarios are:

- (I) Three continuous variables (y_1, y_2, y_3) defined as $y_i = z_i$, for $i = 1, 2, 3$.
- (II) Two binary variables (y_1, y_3) defined as $y_1 = \mathbb{1}(z_1 > 5)$ and $y_3 = \mathbb{1}(z_3 > 3)$ and a continuous variable defined as $y_2 = z_2$.
- (III) Two binary variables (y_1, y_3) defined as in Scenario (II), one ordinal variable y_2 such that $y_2 = \mathbb{1}(4 < z_2 \leq 5) + 2\mathbb{1}(z_2 > 5)$, and a continuous variable $y_4 \sim N(0, 1)$.

The function $\mathbb{1}(\cdot)$ denotes the indicator function that takes the value one if the condition inside is satisfied and zero otherwise.

Scenario I takes the latent continuous vectors as the observed data, so it is expected that the model would not have any problem in detecting the true clustering structure. Scenario II, however, considers only two binary variables defined in terms of the latents z_1 and z_3 , respectively. If we look at Figure 1 we note that binary variables $I(z_1 > 5)$ and $I(z_3 > 3)$ are enough to distinguish the three groups, so the idea with this Scenario II is to test

the definition of the continuous underlying latent variables through thresholding. Finally, Scenario III is the most challenging one because it adds to Scenario II two more variables: one ordinal, y_2 based on z_3 ; and one continuous, y_4 taken from a standard normal. These two additional variables act as noisy variables since the clustering is already defined by the binary variables y_1 and y_3 .

To implement our model, the cut-off points were defined as mentioned in Section 2, that is, $(\gamma_0, \gamma_1, \gamma_2) = (-\infty, 0, \infty)$ for the binary variables, and $(\gamma_0, \gamma_1, \gamma_2, \gamma_3) = (-\infty, 0, 4, \infty)$ for the ordinal variable with 3 categories.

The priors for the parameters a and b that define the Poisson-Dirichlet process were: $\alpha = 0.5$, $d_0^a = d_1^a = d_0^b = d_1^b = 1$. It is well known that the clustering properties of the Bayesian nonparametric mixture models highly rely on the variances (e.g. Barrios et al., 2013; Nieto-Barajas and Contreras-Cristán, 2014). Therefore, we assess the performance of the model by considering different choices for the parameters of the priors on the variances σ_j^2 and $\sigma_{\mu j}^2$. These choices are:

a) $d_0^z = d_0^\mu = 0.1$ and $d_1^z = d_1^\mu = 0.1$

b) $d_0^z = d_0^\mu = 1$ and $d_1^z = d_1^\mu = 1$

c) $d_0^z = d_0^\mu = 2.1$ and $d_1^z = d_1^\mu = 30$

Both prior specifications (a) and (b) are vague and induce a prior variance that goes to infinity. Between the two, prior (a) is the more vague, almost non informative. On the other hand, specification (c) is slightly informative so to induce a prior mean of 27 and a prior variance of 7438. Although the latter specification is slightly more informative than the first two, it can also be considered vague since its variance is still very large.

Combining Scenarios I–III and prior specifications (a)–(c) we have a total of 9 runs. For each of them a sample of size $n = 100$ was taken and a Gibbs sampler with 2200 iterations

was implemented with a burn-in of 200 and a thinning of 2. A total of 1000 MCMC draws were kept for inference.

We summarised the clustering performance by computing the probability density of the number of groups obtained by the model. Results are reported in Figures 2–3. The former contains of the number of clusters along the MCMC for all the 9 runs considered. As we can see the choice of the prior parameters on the variances is crucial for the performance of the model. The number of clusters reduces when we go from prior (a) to (c) (columns one to three in Figure 2, respectively). The reduction is more drastic in presence of continuous variables like in Scenarios I (first row) and Scenario III (third row). To explain why this happens consider Scenario I that considers three continuous variables. Although in this case the latents correspond to the response, when we use the very vague prior (a) the model tends to over fit the data and thus selecting a different μ_i value for each of the 100 data points. The prior choice (b) reduces the number of groups from 100 to around 33, but still not enough. So, in order to recover the three real groups, we need to specify a slightly more informative prior as the one given in (c), where the histograms shows a distinctive mode in 3 groups. Interestingly, the clustering produced when considering only binary variables, like in Scenario II (second row in Figure 2), is less sensible to the choice of the prior on the variances. This is due to the fact that the corresponding latent variables have fixed variance one, so the prior only applies to the variance of the μ_i 's. The mode in the number of groups is 7, 6 and 4 for priors (a), (b) and (c), respectively. Although the mode is in 4 for prior (c), the best clustering, obtained as described in Section 4, has 3 groups as desired.

In Scenario III we have the same two binary variables as in Scenario II plus two more variables, one binary variable that is slightly informative of the clustering and one continuous that is completely unrelated to the underlying clustering. In this scenario the reduction in the number of groups goes from 82 to 6 when moving from prior (a) to (c). Even in the case of prior (c), the model is unable to capture the 3 groups. The best clustering in this case

has 5 groups with 3 large groups and two small groups, which is reasonable good.

Finally, we want to show the evolution in the number of groups inferred by the model along the MCMC iterations. Considering Scenario III with prior (c), Figure 3 presents the number of groups for iterations 1, 11, 20 and 30 with zero burn in and no thinning. This graph shows that the speed in detecting the “correct” number of clusters is very fast, so there is no need for huge chains if we want to explore the clustering space.

As mentioned above, no pre-specified value for a and b were taken. These parameters are also very important in the determination of the number of groups. Their estimated values (posterior means) for scenario I and priors (a), (b) and (c) were 0.99, 0.57 and 0.03 for a , and 0.05, 1.29 and 0.52 for b , respectively. It is interesting to see that the estimated values for a get smaller, inducing a smaller number of groups, as the prior variances become slightly more informative.

5.2 Households data

The ENIGH-MCS is an annual nationwide survey in Mexico that is representative to all the 32 states that form the country. For illustration purposes we concentrate in the year 2014 and in one of these states, the “State of Mexico”. From now on we will refer to it as Edomex for its initials in Spanish. In 2014 this state had 16.2 million inhabitants, that corresponds to 13.5% of the country population, and had 4.2 million households. Edomex is the most populated state of the country. The survey in Edomex consisted of $n = 1,730$ households, which correspond to 0.04% of the total number of households in the state. Since the survey is based on a complex sampling design, each of the households in the sample represents between 960 and 5,286 households with the mode around 2,500 households. A probability histogram of the sampling weights is shown in Figure 4.

The analysis will be based on a vector of $p = 9$ variables of mixed mode. One continuous variable: $Y_1 =$ income (in Mexican pesos); six binary variables: $Y_2 =$ deprivation to feeding

(1–yes, 0–no), Y_3 = deprivation to health services (1–yes, 0–no), Y_4 = housing quality (1–bad, 0–good), Y_5 = education backwardness (1–yes, 0–no), Y_6 = deprivation to basic public services (1–yes, 0–no), Y_7 = deprivation to social security (1–yes, 0–no); one ordinal variable: Y_8 = education level of the family head (0–incomplete primary, 1–incomplete secondary, 2–complete secondary or more); and one nominal variable: Y_9 = town size (1–[100000, ∞), 2–[15000, 100000), 3–[2500, 15000), 4–(0, 2500) inhabitants).

Although variable Y_9 can be treated as ordinal, we prefer to treat it as nominal due to the following reasons. The town sizes intervals are unevenly distributed, and moreover, the Ministry of social development believes that poverty conditions are very different in each of these four strata and typically carries out separated analysis for each category.

According to the procedure described in Section 2, our $p = 9$ dimensional observed vector $\mathbf{Y}$ is associated to a latent vector $\mathbf{Z}$ of dimension $q = 11$. This is the result of associating one latent variable to each of the continuous, binary and ordinal variables plus three latent variables for the nominal variable that has four categories.

For the continuous variable Y_1 we require a normalizing transformation to define Z_1 . An option is to consider the general Box-Cox class with a shift (Box and Cox, 1964) and find the best transformation using an optimality criterion. For the case of income variables we suggest to consider a logarithmic transformation with a shift to avoid having problems for the values near zero, that is $Z_1 = \log(Y_1 + \xi_s)$ with ξ_s the quantile of order s of Y_1 . Specifically we took $s = 0.01$. For the binary variables Y_j , $j = 2, \dots, 7$, the latent variables Z_j have associated thresholds $\gamma_{j,0} = -\infty$, $\gamma_{j,1} = 0$ and $\gamma_{j,2} = \infty$, respectively. For the ordinal variable Y_8 with $K = 3$ ordered categories we define the thresholds as $\gamma_{8,0} = -\infty$, $\gamma_{8,1} = 0$, $\gamma_{8,2} = 4$ and $\gamma_{8,3} = \infty$.

The prior specification for the parameters of the Poisson-Dirichlet process were the same as those taken in the simulation study, i.e., $\alpha = 0.5$, $d_0^a = d_1^a = d_0^b = d_1^b = 1$. To specify the prior on the variances we considered the slightly informative case that arises when $d_0^z =$

$d_0^\mu = 2.1$ and $d_1^z = d_1^\mu = 30$.

For comparison purposes we consider three cases that are described as follows. Let $\bar{w} = (1/n) \sum_{i=1}^n w_i$ then

- i) Ignoring the sampling design, $\pi_i = 1$ and $\kappa = 1$
- ii) Acknowledging the sample design, $\pi_i = 1/w_i$ and $\kappa = 2\bar{w}$
- iii) Acknowledging the sample design, $\pi_i = 1/w_i$ and $\kappa = 4\bar{w}$

We recall that model (1) includes a parameter κ that acts multiplicative in the variance together with the sampling probabilities π_i . As was shown in the simulation study, the variances play a crucial role in determining the clustering structure. Now, since the sampling weights range from 960 to 5286, the sampling probabilities range from 0.00019 to 0.00104. Therefore, in order to prevent having a large number of groups, we use the value of κ to increase the value of the sampling probabilities but keeping the relative importance of each observation. On the other hand, since $\sum_{i=1}^n w_i = \sum_{i=1}^n (1/\pi_i)$ corresponds to the population size, a value of $\kappa \neq 1$ induces a hypothetical population with smaller or larger sizes for $\kappa > 1$ or $\kappa < 1$, respectively.

The Gibbs sampler was run for 2,200 iterations with a burn in of 200 and a thinning of 2. Results for each of the three cases (i)–(iii) are summarised as heat maps of the average identity (clustering) matrix in Figure 5. Ignoring the sampling design, case (i), has the consequence of our model producing 178 groups. On the other hand, when considering the sampling design and for the two different values of κ we obtain 44 groups in case (ii) and 11 groups in case (iii). Although the number of groups obtained from case (iii) is smaller, the corresponding heat map shows that the groups are less homogeneous as compared to the heat maps for cases (i) and (ii).

As it is well known in applied cluster analysis, the “best clustering” does not exist, there are different alternative clusterings and we must choose one according to different criteria

including that of a good interpretation of the groups.

We now proceed to give a brief interpretation of the groups obtained. In Tables 1 and 2 we display the groups (weighted) means for the first eight variables and the (weighted) percentage of households in each of the four categories of the ninth variable. We also include the group size in percentage in the last column. In the last row of the tables we include the population (weighted) means in the whole Edomex for reference. Note that the last value of the table corresponds to the total number of households in Edomex (population size). To simplify the interpretation of the groups we also highlighted in bold those numbers that correspond to a considerably worse poverty condition as compared with the population mean.

Considering first the clustering of 11 groups, we can say the following. Group 8 is the largest group with 39.1% of the households in Edomex and its poverty variables show that it is an average group with no particular deprivation different from the state mean. Group 5 has the largest income and the smallest deprivation indicators which make it to be the wealthier group and represents the 0.5% of the households. The groups 9, 10 and 11 are the poorer groups, with the majority of their values highlighted in bold. Specifically, groups 9 and 11 have the smallest income, but the three groups together consist of households in rural and semi-rural areas (less than 15000 inhabitants) and represent about 10% of the households in the state. Surprisingly the income of group 10 is not that low and the education level of the head of the family is the highest recorded, but still it is a group with health, housing, education, public services and social security deprivations. Group 1 presents an interesting feature, it is formed by households from metropolitan areas (more than 100000 inhabitants) and by households from rural areas (less than 2500 inhabitants), but all present feeding, education and public services deprivations as well as bad education level of the head of the family.

A clustering with 11 groups is very concise and allows us to identify big chunks of house-

holds with certain needs. However a clustering with more groups would allow us to identify households with hidden needs. The clustering of 44 groups might sound too large at a first glance but if we consider that the number of households in the state is around 4 million, then a clustering with 44 groups might not be that large.

Analysing Table 2, more specific needs can be detected in the groups. For instance groups 12 and 42 consists of households in urban areas (more than 15000 inhabitants) but have feeding problems and no access to social security. These two groups are hidden in the thicker clustering of Table 1. Group 43, on the other hand, has the largest income, live in semi-urban areas, but show problems related to health with no access to health services and no social security. Another important aspect of this thinner clustering is that there are no large groups, the group with most households has 8.93% of the total (group 21), whereas the smallest group has 0.04% of the households (group 41). This is in contrast to the clustering with 11 groups where the 85% of the households concentrate in 3 very large groups.

6 Concluding remarks

Multivariate cluster analysis is one of the most useful techniques in practice and there are many methods available in all statistical software. This techniques typically require the data to be numeric and preferably continuous. However, when the multivariate data contains mixed mode variables like continuous, discrete, ordinal and nominal, the clustering task cannot be performed easily.

We discussed a nonparametric model-based clustering approach that is entirely flexible and able to perform clustering of individuals with multivariate mixed-scale variables. The model relies on the introduction of latent continuous variables modelled via multivariate mixture model. The mixing distribution is kept unspecified and totally flexible assuming a nonparametric Poisson-Dirichlet process prior. In addition to the extreme flexibility of the latter approach, its almost sure discreteness is extremely useful to induce the desired

clustering structure.

Our model allows for the treatment of data coming from a complex design, as in our motivating application about Mexican households survey, including the sampling weights in the analysis.

Our findings suggest the importance of the cluster component variances in the clustering procedure and specifically the lower the variances, the higher the number of groups and viceversa. Thus, according to the desired needs, the prior distributions for the variances and any scaling factor need to be carefully elicited.

As for the particular application, our clustering tool will be very beneficial for the authorities in charge of creating social programmes as it allows to identify those households with specific needs and to see the potential number of households that would prosper with the implementation of a social program.

Acknowledgements

The second author acknowledges support from *Asociación Mexicana de Cultura, A. C.* Mexico.

References

- Bandyopadhyay, D. and Canale, A. (2016). Non-parametric spatial models for clustered ordered periodontal data. *Journal of the Royal Statistical Society, Series C* **65**, 619–640.
- Banfield, J.D. and Raftery, A.E. (1993). Model-based Gaussian and non-Gaussian clustering. *Biometrics* **49**, 803–821.
- Barnard, J., McCulloch, R. and Meng, X.-L. (2000). Modeling covariance matrices in terms of standard deviations and correlations, with application to shrinkage. *Statistica Sinica* **10**, 1281–1311.

- Barrios, E., Lijoi, A., Nieto-Barajas, L.E. and Prünster, I. (2013). Modeling with normalized random measure mixture models. *Statistical Science* **28**, 313–334.
- Box, G. E. P. and Cox, D. R. (1964). An analysis of transformations (with discussion). *Journal of the Royal Statistical Society, Series B* **26**, 211–252.
- Canale, A., Dunson, D. B., (2015). Bayesian multivariate mixed-scale density estimation. *Statistics and Its Interface* **8**, 195–201.
- Canale, A. and Scarpa, B. (2016). Bayesian nonparametric location-scale-shape mixtures. *Test* **25**, 113–130.
- Chambers, R. L. and Skinner, C. J. (2003). *Analysis of Survey Data*. Wiley, Chichester.
- Coneval (2009). *Metodología para la medición multidimensional de la pobreza en México*. Consejo Nacional de Evaluación de la Política de Desarrollo Social, México. (In Spanish). Available at: http://www.coneval.org.mx/rw/resource/Metodologia_Medicion_Multidimensional.pdf
- Dahl, D.B. (2006). Model based clustering for expression data via a Dirichlet process mixture model. In *Bayesian Inference for Gene Expression and Proteomics*, Eds. M. Vanucci, K.-A. Do and P. Müller. Cambridge University Press, Cambridge.
- Everitt, B.S. (1988). A finite mixture models for the clustering of mixed-mode data. *Statistics and Probability Letters* **6**, 305–309.
- Ferguson, T.S. (1973). A Bayesian analysis of some nonparametric problems. *Annals of Statistics* **1**, 209–230.
- Ishwaran, H. and James, L.F. (2001). Gibbs sampling methods for stick-breaking priors. *Journal of the American Statistical Association* **96**, 161–173.

- Kingman, J.F.C. (1975). Random discrete distributions. *Journal of the Royal Statistical Society, Series B* **37**, 1–22.
- Kottas, A., Müller, P. and Quintana, F. (2005). Nonparametric Bayesian modeling for multivariate ordinal data. *Journal of Computational and Graphical Statistics* **14**, 610–625.
- Leon-Novelo, L.G., Zhou, X., Nebiyu Bekele, B. and Müller, P. (2010). Assessing toxicities in a clinical trial: Bayesian inference for ordinal data nested within categories. *Biometrics* **66**, 966–974.
- Lumley, T. (2010). *Complex Surveys*. Wiley, New Jersey.
- McParland, D., Claire Gormley, I., McCormick, T.H., Clark, S.J., Whiteson Kabudula, C. and Collinson, M.A. (2014). Clustering South African households based on their asset status using latent variable models. *Annals of Applied Statistics* **8**, 747–776.
- Navarrete, C., Quintana, F.A. and Müller, P. (2008). Some issues in nonparametric Bayesian modeling using species sampling models. *Statistical Modelling* **8**, 3–21.
- Nieto-Barajas, L.E. and Contreras-Cristán, A. (2014). A Bayesian nonparametric approach for time series clustering. *Bayesian Analysis* **9**, 147–170.
- Norets, A., Pelenis, J., (2012). Bayesian modeling of joint and conditional distributions. *Journal of Econometrics* **168**, 332–346.
- Pitman, J. (1995). Exchangeable and partially exchangeable random partitions. *Probability Theory and Related Fields* **102**, 145–158.
- Pitman, J. and Yor, M. (1997). The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator. *The Annals of Probability* **25**, 855–900.

- R Core Team (2016). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Ritter, C. and Tanner, M. A. (1992). Facilitating the Gibbs sampler. The Gibbs stopper and the Griddy-Gibbs sampler. *Journal of the American Statistical Association* **87**, 861–868.
- Rodríguez, C. E. and Walker, S. G. (2014). Univariate Bayesian nonparametric mixture modeling with unimodal kernels. *Statistics and Computing* **24**, 35–49.
- Tierney, L. (1994). Markov chains for exploring posterior distributions. *Annals of Statistics* **22**, 1701-1722.
- Wade, S. and Ghahramani, Z. (2015). Bayesian cluster analysis: Point estimation and credible balls. Preprint.

Table 1: Group means for the clustering with 11 groups obtained with case (iii). Columns are divided according to the nature of the variables: continuous, binary, ordinal and nominal. Last row is the population mean, apart from the last value that corresponds to the population size.

group	income	feed	health	house	edu	serv	ss	hedu	ts:1	ts:2	ts:3	ts:4	group.s
1	2081	0.31	0.19	0.07	1.00	0.44	0.55	0.25	0.36	0.05	0.19	0.39	4.15%
2	5819	0.31	0.35	0.12	0.95	0.15	0.62	0.37	0.54	0.07	0.27	0.13	2.81%
3	4739	0.21	0.36	0.11	0.37	0.14	0.83	1.87	0.53	0.15	0.16	0.16	28.06%
4	44128	0.04	0.11	0.00	0.16	0.02	0.30	2.00	0.65	0.29	0.02	0.04	2.91%
5	219578	0.16	0.27	0.00	0.00	0.00	0.27	2.00	0.89	0.11	0.00	0.00	0.46%
6	20296	0.09	0.36	0.03	0.27	0.05	0.70	1.99	0.72	0.11	0.12	0.05	17.77%
7	2235	0.37	0.51	0.14	0.50	0.18	0.93	1.61	0.53	0.06	0.22	0.20	3.88%
8	9284	0.19	0.33	0.08	0.33	0.08	0.72	1.92	0.65	0.11	0.14	0.10	39.13%
9	425	0.52	0.23	0.23	0.69	0.54	1.00	1.11	0.23	0.16	0.24	0.37	0.72%
10	6551	0.00	1.00	1.00	1.00	1.00	1.00	2.00	0.00	0.00	0.00	1.00	0.06%
11	805	1.00	1.00	1.00	1.00	0.00	1.00	1.00	0.00	0.00	1.00	0.00	0.05%
pop	11212	0.19	0.34	0.08	0.38	0.11	0.74	1.79	0.61	0.12	0.15	0.13	4240837

Table 2: Group means for the clustering with 44 groups obtained with case (ii). Columns are divided according to the nature of the variables: continuous, binary, ordinal and nominal. Last row is the population mean, apart from the last value that corresponds to the population size.

group	income	feed	health	house	edu	serv	ss	hedu	ts:1	ts:2	ts:3	ts:4	group.s
1	4912	0.11	0.86	0.07	0.24	0.05	1.00	1.97	0.68	0.10	0.13	0.10	5.40%
2	6529	0.24	0.39	0.36	0.63	0.98	0.95	1.91	0.09	0.17	0.35	0.40	1.26%
3	11091	0.19	0.08	0.14	0.92	0.08	0.88	1.88	0.54	0.15	0.16	0.14	4.74%
4	185628	0.18	0.18	0.00	0.00	0.00	0.18	2.00	1.00	0.00	0.00	0.00	0.41%
5	4610	0.12	0.27	0.97	0.24	0.17	0.72	1.95	0.50	0.29	0.08	0.13	1.31%
6	4854	0.06	0.07	0.06	0.17	0.06	0.14	1.90	0.52	0.25	0.10	0.14	2.63%
7	23345	0.02	0.29	0.02	0.13	0.00	0.94	1.99	0.81	0.10	0.08	0.01	5.24%
8	4830	0.10	0.15	0.00	0.89	0.11	0.95	1.89	0.47	0.10	0.27	0.16	2.18%
9	8908	0.17	0.91	0.05	0.09	0.03	0.98	1.97	0.71	0.12	0.13	0.05	6.23%
10	48232	0.00	0.03	0.00	0.10	0.00	0.20	2.00	0.60	0.40	0.00	0.00	2.09%
11	3629	0.08	0.74	0.12	0.83	0.87	1.00	1.79	0.13	0.00	0.24	0.63	1.30%
12	6282	0.97	0.16	0.09	0.14	0.05	0.91	1.93	0.69	0.10	0.10	0.11	3.94%
13	8171	0.08	0.88	0.10	0.90	0.13	0.96	1.88	0.63	0.10	0.13	0.15	4.86%
14	24776	0.00	0.00	0.00	0.05	0.01	0.06	2.00	0.84	0.06	0.07	0.03	3.53%
15	8496	0.10	0.01	0.04	0.16	0.04	0.19	1.97	0.71	0.10	0.13	0.06	7.77%
16	4233	0.20	0.18	0.16	0.96	0.18	0.76	0.69	0.39	0.10	0.24	0.28	2.75%
17	2253	0.23	0.87	0.04	0.09	0.15	1.00	1.80	0.69	0.07	0.13	0.10	1.61%
18	11271	0.07	0.10	0.01	0.02	0.01	0.91	1.98	0.59	0.13	0.20	0.08	5.57%
19	3238	0.85	0.60	0.12	0.94	0.23	0.95	1.94	0.40	0.16	0.28	0.16	2.16%
20	14557	0.07	0.06	0.01	0.11	0.03	0.08	2.00	0.71	0.12	0.12	0.06	4.85%
21	6388	0.04	0.08	0.05	0.08	0.08	0.88	1.93	0.64	0.11	0.14	0.11	8.93%
22	9657	0.92	0.89	0.20	0.92	0.11	0.92	1.96	0.58	0.26	0.08	0.09	1.31%
23	3173	0.26	0.08	0.03	0.23	0.10	0.87	1.91	0.59	0.15	0.06	0.19	4.78%
24	6294	0.44	0.82	0.11	0.96	0.17	1.00	0.60	0.46	0.15	0.28	0.11	1.59%
25	8054	0.18	0.00	0.00	0.86	0.00	0.14	0.60	0.79	0.00	0.14	0.07	0.82%
26	3204	0.00	0.05	0.00	1.00	0.00	0.05	0.22	0.65	0.00	0.29	0.06	1.12%
27	17282	0.12	0.79	0.04	0.45	0.11	0.99	1.97	0.66	0.10	0.12	0.12	5.37%
28	31499	0.00	1.00	0.07	0.65	0.14	1.00	2.00	0.55	0.09	0.29	0.08	0.75%
29	2159	0.58	0.05	0.17	1.00	0.96	0.70	0.23	0.04	0.10	0.18	0.68	1.21%
30	408	0.87	0.12	0.00	0.62	0.37	1.00	1.62	0.24	0.11	0.41	0.24	0.43%
31	1455	0.34	0.36	0.08	1.00	0.27	0.70	0.21	0.37	0.05	0.23	0.35	1.88%
32	21621	1.00	0.28	0.00	0.90	0.00	0.71	1.90	0.82	0.00	0.18	0.00	0.62%
33	1273	0.55	0.58	0.30	0.84	0.42	1.00	1.59	0.16	0.00	0.26	0.58	0.38%
34	4136	0.00	1.00	1.00	1.00	0.00	1.00	1.00	1.00	0.00	0.00	0.00	0.05%
35	458	0.00	0.30	0.32	1.00	1.00	1.00	0.00	0.00	0.39	0.00	0.61	0.18%
36	695	0.51	0.25	1.00	1.00	0.23	1.00	1.26	0.00	0.00	0.77	0.23	0.21%
37	4243	0.00	1.00	1.00	0.00	1.00	1.00	2.00	0.00	0.00	1.00	0.00	0.06%
38	5390	0.00	0.00	0.48	1.00	1.00	0.48	0.48	0.48	0.00	0.52	0.00	0.11%
39	4498	0.51	0.00	1.00	1.00	1.00	0.00	0.00	0.00	0.00	0.51	0.49	0.10%
40	17110	1.00	0.00	0.00	1.00	0.00	0.00	0.00	1.00	0.00	0.00	0.00	0.05%
41	569	0.00	0.00	1.00	1.00	0.00	1.00	1.00	1.00	0.00	0.00	0.00	0.04%
42	2259	1.00	0.00	0.00	0.00	0.00	1.00	1.00	1.00	0.00	0.00	0.00	0.06%
43	499434	0.00	1.00	0.00	0.00	0.00	1.00	2.00	0.00	1.00	0.00	0.00	0.05%
44	51837	1.00	0.00	0.00	1.00	1.00	1.00	2.00	0.00	0.00	1.00	0.00	0.06%
pop	11212	0.19	0.34	0.08	0.38	0.11	0.74	1.79	0.61	0.12	0.15	0.13	4240837




Figure 1: Simulated data set over the three latent variables Z_1, Z_2, Z_3 .




Figure 2: Histograms of the number of clusters for scenarios I, II and III (in the rows) and prior specifications (a), (b) and (c) in the columns.




Figure 3: Evolution of the model clustering process.




Figure 4: Probability histogram of the sampling design weights or expansion factors w_i .




Figure 5: Heatmaps representation of average clustering matrix for different prior specifications. Ignoring sample weights and $\kappa = 1$ (top left), with sample weights and $\kappa = 2\bar{w}$ (top right), and $\kappa = 4\bar{w}$ (bottom).